



Estimated Vegetated Cover in 50 Meter Stream and Lake Buffer

This EnviroAtlas community map portrays the estimated percent of vegetated cover within 50 meters of streams, rivers, and other hydrologically connected waterbodies (e.g., lakes and ponds) for each census block group.

Why are stream buffers important?

Vegetation adjacent to streams and rivers, also called the [riparian](#) area (or [riparian buffer](#)), helps protect terrestrial wildlife habitat, aquatic habitat, and water quality. Natural riparian buffers may exist at varying distances from a waterbody depending on the occurrence and width of flood plains, hydric soils, or associated wetlands. In more developed areas, a riparian buffer represents a vegetated area occurring between a waterbody and adjacent human disturbances. Maintaining vegetated cover in stream buffers benefits water quality at the site as well as downstream. Land management in upstream areas directly affects the water quality in downstream rivers, bays, and estuaries.

By slowing and processing stormwater runoff, riparian vegetation can prevent sediment, nutrients, harmful bacteria, pesticides, and metals from entering the waterway and degrading water quality. By storing water and securing the stream bank with roots, riparian trees can reduce flooding, help prevent erosion, and minimize downstream property damage. The ability of riparian vegetation to filter pollutants and store floodwater varies with local climate, buffer width, slope, soil permeability, and depth to water table.¹

Studies have shown that sediment removal by trees ranges from 60–90% depending on buffer area, slope, and the volume and velocity of runoff.¹ Toxic substances adhering to sediment particles may be modified by soil microorganisms into less harmful forms and made available to plants. A published review of 66 studies covering nutrient removal by mixed buffer vegetation found that 75% and 90% of excess nitrogen was removed from mean buffer widths of 28 and 112 meters (92 and 367 feet), respectively.²

Vegetated stream buffers provide critical wildlife habitat for resident and migratory wildlife species that depend on riparian areas for cover, food, and water. Buffers create corridors for wildlife to safely move between patches of habitat. Riparian vegetation provides canopy cover for the waterbody, helping to regulate the temperature of the water and protect habitat for fish, amphibians, and aquatic insects. The benefits of aquatic and riparian habitat extend beyond




Photo: Sacramento Riparian Buffer, USFWS

the stream segment to support biodiversity downstream, throughout the watershed, and in the surrounding region.

Although the ecosystem services provided by riparian buffers are well-known, determining the optimum widths necessary for buffers to deliver specific benefits and functions (e.g., flood storage, temperature moderation, nutrient filtering) is more difficult. Streams with adjacent intense disturbances require wider buffers.³ Narrow buffer widths of 5–15 meters (16–49 feet) maintain bank stability and provide some temperature moderation, but they are inadequate for sediment and nutrient reduction.³ Narrow buffer strips are also subject to flood and wind damage. Maintaining breeding habitat for songbirds and wildlife corridors for the movement of large mammals requires wider buffer widths of 30.5–91.4 meters (100–300 feet).^{3,4}

How can I use this information?

The map, Estimated Vegetated Cover in 50 Meter Stream and Lake Buffer, is one of six EnviroAtlas maps that illustrate land area, tree cover, and vegetated cover within 50 meters of the banks of community streams and lakes. The related map, Percent Vegetated Cover in 50 Meter Stream and Lake Buffer, can be overlaid on this layer to summarize the vegetated cover in buffer by census block group. An area can be more thoroughly investigated by increasing the transparency of this map and adding layers for streams and waterbodies (NHDPlus, found under the boundaries icon) to an aerial imagery base map. Other background layers are available, such as community land cover maps created for EnviroAtlas at 1-meter resolution (enter *land cover*

community in the interactive map search box) and community boundaries (under the boundaries icon). Detailed examination shows land cover along streams and reveals where upstream areas may be contributing to water quality problems downstream. Map analysis can help identify disparities in benefits among block groups, for example, flood-prone areas or source areas of heavy metals or nutrients that lack nearby woodland to filter pollutants. Planners and other interested users can explore where riparian restoration would have the greatest return in terms of improving water quality in nearby waterbodies. Many states have developed guidelines for riparian buffer best management practices (BMPs) and may include recommended buffer widths for various functions.

How were the data for this map created?

This map is based on the land cover data derived for each EnviroAtlas community and the high resolution (scale 1:24,000 or higher) National Hydrography Dataset Plus Version 2 ([NHDPlusV2](#)). The land cover data were created from one-meter aerial photography through remote-sensing methods. Vegetated cover includes trees, forests, grass, herbaceous cover, shrubs, and woody and emergent wetlands, but it excludes agriculture and orchards because of the potential application of pesticides and fertilizers that could pollute waterways. Two moving window analyses and basic raster math first determined the percent of vegetated cover within 50 meters of every point in the community. Then, the NHD stream and lake edges were used to extract only those windows that captured percent vegetated cover in the center of each 50 meter buffer, representing the width of the buffer. Waterbodies that are not hydrologically connected were not included in the analysis.

What are the limitations of these data?

All of the EnviroAtlas community maps that are based on land cover rely on aerial imagery and other remotely-sensed data. These data are estimates that are inherently imperfect. The land cover maps used in the community component of EnviroAtlas typically have an overall accuracy of between

80 and 90%, meaning that there is a probability of at least 80 percent that the land cover reported at any given point corresponds to objects in the native aerial photography. The quality of the NHD streamlines varies between regions, and the data may not capture all streams. These data will be updated over time; updates are expected to improve accuracy as data and classification methods improve.

Although fixed-distance buffers are commonly used in spatial analysis, they do have limitations relative to variable-distance buffers. Along a stream network, riparian cover may be narrower or wider than the fixed-distance buffer. Fixed-distance buffers cannot account for differences among buffer areas because of gaps in riparian vegetation, upslope sources of pollutants, or upslope forested areas.⁵ They do not reflect upstream-downstream patterns of watershed land cover, differences between forested and unforested stream banks, or flowpaths for runoff influenced by local topography.⁵ A full research effort, one that considered variable buffer widths, would be required to get a more accurate estimate of local buffer filtering capabilities.

How can I access these data?

EnviroAtlas data can be viewed in the interactive map, accessed through web services, or downloaded.

Where can I get more information?

A selection of resources related to riparian buffers and their benefits is listed below. For additional information on data creation, access the metadata found in the drop-down menu for each map layer. To ask specific questions about this data layer, please contact the [EnviroAtlas Team](#).

Acknowledgments

EnviroAtlas is a collaborative effort by the US EPA, its contractors, and project partners. Alexandra Sears, Student Services Contractor, developed this map. The fact sheet was created by Anne Neale and Donald Ebert, EPA, Alexandra Sears and Jessica Jahre, Student Services Contractors, and Sandra Bryce, Innovate!, Inc.

Selected Publications

1. Nowak, D.J., J. Wang, and T. Endreny. 2007. [Chapter 4: Environmental and economic benefits of preserving forests within urban areas: air and water quality](#). Pages 28–47 in de Brun, C.T.F. (ed.), *The economic benefits of land conservation*. The Trust for Public Land, San Francisco, California.
2. Mayer, P.M., S.K. Reynolds, M.D. McCutchen, and T.J. Canfield. 2006. [Riparian buffer width, vegetative cover, and nitrogen removal effectiveness: A review of current science and regulations](#). EPA/600/R-05/118. U.S. Environmental Protection Agency, Cincinnati, Ohio.
3. Palone, R.S., and A.H. Todd (eds.). 1997. [Chesapeake Bay riparian handbook: A guide for establishing and maintaining riparian forest buffers](#). NA-TP-02-97, U.S. Forest Service, Radnor, Pennsylvania.
4. Bentrup, G. 2008. [Conservation buffers: Design guidelines for buffers, corridors, and greenways](#). General Technical Report SRS-109, U.S. Forest Service, Southern Research Station, Asheville, North Carolina. 110 p.
5. Baker, M.E., D.E. Weller, and T.E. Jordan. 2006. [Improved methods for quantifying potential nutrient interception by riparian buffers](#). *Landscape Ecology* 21 (8): 1327–1345.