



NIB Small Game Species Richness: Southwest

Desert cottontail under a desert shrub. Photo by G.M. Stolz, USFWS. This EnviroAtlas national map displays the Normalized Index of Biodiversity (NIB), an index value for small game species richness based on potential habitat within each 12-digit hydrologic unit (HUC) in the southwestern United States (Arizona, Colorado, Nevada, New Mexico, and Utah). Small game refers to smaller upland animals that are hunted; their designation as a game species is determined by state fish and wildlife agencies. These data are based on habitat models, not wildlife counts. Potential habitat may be specific to wintering, breeding, or year-round activities depending on the species.



Why are small game species important?

Small game species vary by state and may include grouse, quail, squirrels, pheasants, doves, prairie dogs, and rabbits. Small game species richness estimates the number of small game species that may inhabit an area, based on potential habitat. Species richness is frequently used as a measure of the relative conservation value of a particular area and as a surrogate for measuring [biodiversity](#).

Many scientists believe biodiversity, as it represents all forms of life on earth, provides or supports the core benefits that humans derive from their environment and helps sustain human culture throughout the world. Therefore, many organizations consider managing areas for biodiversity as a means to achieve an acceptable balance among competing demands for various ecosystem services. Small game species richness is one indicator of biodiversity within an area.

Each species plays an important role within its ecosystem and the [food web](#). Within a [food chain](#), small game animals are [primary consumers](#) that serve as an important food source for other wildlife ([secondary consumers](#)). Prairie dogs, for example, are considered a [keystone species](#) for their capacity to maintain grassland areas, serve as prey, and support numerous other species in their system of tunnels. Population declines in some typically abundant small game species, such as the sage-grouse, can indicate an imbalance in their ecosystem.

In addition to the roles that game species play in [ecosystems](#), they are also an important food source and are appreciated for the recreation opportunities and the aesthetic value they provide. Hunting for animals such as pheasants and rabbits

has a long tradition in the U.S.; in 2011, hunters spent \$2.6 billion on hunting small game species.¹

Beyond its recreational value, hunting provides a critical economic vehicle for conservation, management, and restoration projects, the benefits of which extend beyond small game species. Revenue is collected from a federal excise tax, as well as from state hunting licenses and permits. These funds are used to support wildlife and conservation programs and allow for the conservation of lands upon which small game species depend. The income generated by the abundance of small game species helps protect land, maintain wildlife populations, and conserve biodiversity.

How can I use this information?

The map, NIB Small Game Species Richness: Southwest, is one of three EnviroAtlas maps that illustrate indicators of small game species richness for the Southwest. Additional EnviroAtlas maps show the maximum and mean small game species richness for each 12-digit HUC. Used together or independently, these maps can help identify areas of potentially low or high small game species richness to help inform decisions about resource restoration, use, and conservation. Knowing small game species richness provides one aspect necessary to conserve biodiversity.

These maps can also be used in conjunction with other maps in EnviroAtlas to help identify areas with high ecological or recreational value for inclusion in conservation or restoration planning or protection from further development for

recreational or aesthetic reasons. This information can help identify areas that may be vulnerable to development.

After finding the small game species richness values for a particular 12-digit HUC, an area can be more intensively investigated by using individual species models at a higher resolution. Individual species models are available through the Southwest Regional Gap Analysis Project ([SWReGAP](#)).

How were the data for this map created?

This data layer is based on data generated by the U.S. Geological Survey (USGS) National Gap Analysis Program ([GAP](#)). The GAP program maps the distribution of natural vegetation communities and potential habitat for individual terrestrial vertebrate species. These models utilize predictive environmental variables (e.g., GAP land cover, elevation, distance to water) to derive deductive habitat models for each species.

A component of GAP, SWReGAP modeled habitat for 36 small game species that reside, breed, or use the habitat within the 5-state southwest study area for a significant portion of their life history. Small game species richness was calculated by combining predicted habitat for all GAP individual small game species by pixel across the southwestern United States. The number of small game species in each pixel was then summarized by 12-digit HUC and the mean value calculated for each HUC. The NIB was calculated by dividing the mean value by the maximum value for each HUC.

What are the limitations of these data?

EnviroAtlas uses the best data available, but there are still limitations associated with the data. These data are based on models and large national geospatial databases. Calculations based on the data are estimations of the truth founded on the best available science. Modeled data can be complementary but the information is not meant to replace monitoring data. Habitat models do not predict the actual occurrence of

species, but rather their predicted occurrence based on their known associations with certain habitat types. Habitat is only one factor that determines the actual presence of a species. Other factors include habitat quality, predators, prey, competing species, and fine scale habitat features such as woody debris.

How can I access these data?

EnviroAtlas data can be viewed in the interactive map, accessed through web services, or downloaded. Metric values for individual pixels may be obtained from the [New Mexico State University Center for Applied Spatial Ecology](#). [SWReGAP](#) and [GAP](#) data and accuracy information can be accessed through their respective websites.

Where can I get more information?

There are numerous resources about the importance of game species and on biodiversity in general; a selection of these resources is below. Information on what is considered small game, and applicable management, rules, and regulations, can be found on each state's fish and wildlife department website. For additional information on how the data were created, access the metadata for the data layer from the drop down menu on the interactive map table of contents and click again on metadata at the bottom of the metadata summary page for more details. To ask specific questions about this data layer, please contact the [EnviroAtlas Team](#).

Acknowledgments

EnviroAtlas is a collaborative effort led by EPA. The data for Small Game Species Richness were created through a collaborative effort between the USGS GAP and EPA. The data were generated by Kenneth Boykin and graduate students from New Mexico State University. The data used to derive Small Game Species Richness came from [SWReGAP](#). The fact sheet was written by Kenneth Boykin, New Mexico State University, Anne Neale and William Kepner, EPA, and Jessica Jahre, EPA Student Services Contractor.

Selected Publications

1. U.S. Department of the Interior, U.S. Fish and Wildlife Service, and U.S. Department of Commerce, U.S. Census Bureau. 2013. [2011 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation](#), FHW/11-NAT (RV), Washington, D.C.
- Boykin, K.G., W.G. Kepner, D.F. Bradford, R.K. Guy, D.A. Kopp, A. Leimer, E. Samson, F. East, A. Neale, and K. Gergely. 2013. [A national approach for mapping and quantifying habitat-based biodiversity metrics across multiple spatial scales](#). *Ecological Indicators* 33:139–147.
- Boykin, K.G., B.C. Thompson and S. Propeck-Gray. 2010. [Accuracy of gap analysis habitat models in predicting physical features for wildlife-habitat associations in the southwest U.S.](#) *Ecological Modelling* 221:2769–2775.
- Kepner, W. G., K. G. Boykin, D. F. Bradford, A. C. Neale, A. K. Leimer, and K. J. Gergely. 2011. [Biodiversity metrics fact sheet](#). U.S. Environmental Protection Agency, Washington, DC, EPA/600/F-11/006.
- Prior-Magee, J.S., K.G. Boykin, D.F. Bradford, W.G. Kepner, J.H. Lowry, D.L. Schrupp, K.A. Thomas, and Bruce C. Thompson, Editors. 2007. [Southwest Regional Gap Analysis Project Final Report](#). U.S. Geological Survey, Gap Analysis Program, Moscow, ID.