



Percent Natural Land Cover in Stream Buffer

This EnviroAtlas national map portrays the percent of land within 45 meters of streams, rivers, and other hydrologically-connected waterbodies (e.g., lakes and ponds) covered by natural land cover within each 12-digit hydrologic unit (HUC). The map uses the 2011 USDA Cropland Data Layer (CDL) combined with the 2011 National Land Cover Database (NLCD) to define natural land cover (forest, shrub, wetland, and grassland/herbaceous). Land covers like agriculture and developed land are excluded.

Why are stream buffers important?

Natural land cover adjacent to streams and rivers, the [riparian area](#) (or [riparian buffer](#)), helps protect terrestrial wildlife habitat, aquatic habitat, and water quality. Riparian areas store stormwater and runoff, filter pollutants from the air and soil, and moderate air and water temperatures. Maintaining natural vegetation in stream buffers benefits water quality at the site as well as downstream. Land management in upstream areas directly affects the water quality in downstream rivers, bays, and estuaries. Natural stream buffers can provide air quality, carbon storage, and climate stabilization benefits while adding recreation, cultural, and aesthetic value to an area. The quantity and quality of benefits provided by riparian buffers depend on local conditions (e.g., climate, slope, buffer width, soil permeability, pollutant load, and depth to water table).^{1,2}

Riparian areas intercept and filter stormwater and runoff. By trapping and processing pollutants, riparian vegetation can prevent sediment, nutrients, harmful bacteria, pesticides, and other pollutants from entering waterbodies. Riparian vegetation can help regulate the flow of flood water into a waterbody by slowing runoff, allowing it to soak into the ground to recharge ground water, and transpiring moisture back into the atmosphere through leaves and stems. Forested riparian buffers provide as much as 40 times the flood water storage of a cropped agricultural field and 15 times that of grass turf because of the ability of tree canopy to capture rainfall on the surface area of the leaves, stems, and branches and to take up water through deep roots.² By storing water and securing the stream bank with roots, riparian vegetation can reduce flood potential, help prevent erosion, and minimize downstream property damage.

Riparian vegetation, particularly tree cover, can regulate the temperature of the water and protect habitat for fish and other aquatic life. Elevated temperatures increase the impacts of nonpoint source pollution and deplete oxygen from the



Photo: Iowa stream, L. Betts, NRCS

system.² Small fish, some amphibians, and most aquatic insects also rely on leaf litter from overhanging vegetation as their primary food source.

Narrow buffer widths of 5–15 meters (16–49 feet) maintain bank stability and provide some temperature moderation, but they are inadequate for sediment and nutrient reduction.² Narrow buffer strips are also subject to flood and wind damage. Streams with adjacent disturbances require wider buffers.² A published review of 66 studies covering nutrient removal by buffer vegetation (grass, grass and trees, forest) found that 75% and 90% of excess nitrogen was removed from mean buffer widths of 28 and 112 meters (92 and 367 feet), respectively.³ Though trees return a significant portion of the nitrogen they remove back to the soil as leaf litter, trees also enable [denitrification](#), a process where bacteria in saturated soil transform dissolved nitrates into gaseous nitrogen compounds that escape to the atmosphere.¹

Natural vegetation in stream buffers provides critical wildlife habitat for resident and migratory species that depend on riparian areas for cover, food, and water. Vegetated buffers create critical corridors for wildlife to safely move under cover from one area to another. Maintaining breeding habitat for songbirds and wildlife corridors for the movement of large mammals requires wider buffer widths of 30.5–91.4 meters (100–300 feet).^{2,4} The benefits of aquatic and riparian habitat extend beyond the stream by maintaining biodiversity downstream and throughout the watershed.

How can I use this information?

This map layer indicates which 12-digit HUCs may benefit from riparian buffer restoration projects to improve water quality. An area can be more thoroughly investigated by increasing the transparency of the map and adding data for streams and water bodies (NHDPlus from Boundary Layers) to an aerial imagery base map. Examination of these layers in more detail shows land cover along streams and reveals where upstream areas may be contributing to problems in downstream communities. Many states have developed guidelines for riparian buffer best management practices (BMPs) and recommended buffer widths.⁴

How were these data created?

These data were generated by using the combined 2011 Cropland Data Layer and 2011 National Land Cover Database (NLCD-CDL) with medium resolution (1:10,000 or higher) [NHDPlusV2](#) National Hydrography Dataset (stream lines and water bodies) in the landscape assessment tool, Analytical Tools Interface for Landscape Assessments (ATtILA). [ATtILA](#) is an Esri ArcGIS 10.0 extension created by EPA that calculates many commonly-used landscape metrics, including land cover adjacent to streams. The 45-meter stream buffers for this group of EnviroAtlas metrics were generated by delineating a polygon 45 meters wide on either side of a stream network and around the perimeter of hydrologically-connected lakes or ponds. The percent natural land cover for the area (NLCD-CDL classified forest, shrubland, grassland, barren land, and wetland) contained within the buffer polygon was recorded for each 12-digit HUC. Waterbodies not hydrologically connected were not included in the analysis. For more information on this calculation, see the [ATtILA User's Manual](#).

What are the limitations of these data?

Though EnviroAtlas uses the best data available, there are limitations associated with the data. The landcover classes found in the NLCD and CDL are created through the classification of satellite imagery. Human classification of

landcover types that have a similar spectral signature can result in classification errors. As a result, NLCD is a best estimate of actual landcover. Also, because of their 30m pixel size, the NLCD and CDL may miss riparian buffers that are <30m wide.

A national-scale metric such as this gives an overview of the extent of natural land cover within a fixed-distance buffer summarized by 12-digit HUCs. However, at any point along a stream network, natural land cover may be narrower or wider than the fixed-distance buffer. Fixed-distance buffers cannot account for differences among buffer areas caused by gaps in riparian vegetation, upslope sources of pollutants, or upslope forested areas.⁵ They do not reflect upstream-downstream patterns of watershed land cover, differences between forested and unforested stream banks, or flowpaths for runoff, influenced by local topography.⁵ A full research effort, one that considered variable buffer widths, would be required to get an accurate local estimate of riparian vegetation filtering capabilities within or among watersheds.

How can I access these data?

EnviroAtlas data can be viewed in the interactive map, accessed through web services, or downloaded. The [NLCD](#), [CDL](#), and [NHDPlusV2](#) data are accessible through their respective websites.

Where can I get more information?

A selection of resources related to riparian buffers is listed below. For additional information on how the data were created, access the [metadata](#) for the data layer. To ask specific questions about this data layer, please contact the [EnviroAtlas Team](#).

Acknowledgments

Donald Ebert, EPA, developed this map for EnviroAtlas. This fact sheet was created by Anne Neale and Donald Ebert, EPA, Jessica Jahre, Student Services Contractor, and Sandra Bryce, Innovate!, Inc.

Selected Publications

1. Nowak, D.J., J. Wang, and T. Endreny. 2007. [Chapter 4: Environmental and economic benefits of preserving forests within urban areas: Air and water quality](#). Pages 28–47 in de Brun, C.T.F. (ed.), *The economic benefits of land conservation*. The Trust for Public Land, San Francisco, California.
2. Palone, R.S., and A.H. Todd (eds.). 1997. [Chesapeake Bay riparian handbook: A guide for establishing and maintaining riparian forest buffers](#). NA-TP-02-97, U.S. Forest Service, Radnor, Pennsylvania.
3. Mayer, P.M., S.K. Reynolds, M.D. McCutchen, and T.J. Canfield. 2006. [Riparian buffer width, vegetative cover, and nitrogen removal effectiveness: A review of current science and regulations](#). EPA/600/R-05/118. U.S. Environmental Protection Agency, Cincinnati, Ohio.
4. Bentrup, G. 2008. [Conservation buffers: Design guidelines for buffers, corridors, and greenways](#). General Technical Report SRS-109. U.S. Forest Service, Southern Research Station, Asheville, North Carolina. 110 p.
5. Baker, M.E., Weller, D.E., Jordan, T.E. 2006. [Improved methods for quantifying potential nutrient interception by riparian buffers](#). *Landscape Ecology*, 21 (8): 1327–1345.