



Percent Tree Canopy Cover

This EnviroAtlas national map estimates the percent of land area that is covered by tree canopy within each 12-digit hydrologic unit ([HUC](#)) in the conterminous U.S using the 2011 National Land Cover Database – [Percent Tree Canopy](#) dataset.

Why is tree canopy cover important?

Percent tree canopy cover records the proportion of land surface that is covered by tree tops. Tree cover helps reduce the potential adverse health and environmental effects of air pollutants and particulates by removing pollutants from the air. Trees can remove significant amounts of air pollution in metropolitan areas. Gaseous air pollutants are taken in primarily through the leaf surface.¹ Trees remove particulates by directly intercepting airborne particles. Some particles can be absorbed into the tissues of the tree, though most particles are retained on the tree surface. Vegetation is a temporary retention site for many atmospheric particles because particulates may be resuspended back to the atmosphere, washed off by rain, or dropped to the ground with leaf and twig fall.²

Tree cover is a major contributor to climate change mitigation. Carbon dioxide is the dominant [greenhouse gas \(GHG\)](#) released into the atmosphere as a result of human activities.³ Trees are composed largely of carbon and they continue to take in carbon as they grow. By fixing carbon during photosynthesis and storing it as biomass, growing trees act as a [sink](#) for CO₂. However, the loss of existing trees from disturbances such as land use change and fires can release this carbon back into the atmosphere, increasing the levels of atmospheric carbon.⁴ Maximizing tree canopy carbon storage is important for maintaining a stable climate.

The proportion of tree canopy cover relative to cropland in agricultural areas and [impervious surfaces](#) in communities influences the quantity, quality, and speed of stormwater runoff entering waterways. Besides providing shade and cooling water temperatures, tree cover in [riparian buffers](#) slows and stores floodwater and filters significant quantities of sediment, nutrients, and heavy metals from agricultural fields and urban runoff. For example, studies have shown that sediment removal by trees ranges from 60–90% depending on buffer area, slope, and the volume and velocity of runoff.⁵ Toxic substances adhering to sediment particles may be modified by soil microorganisms into less harmful forms and made available to plants.



Photo: Mt. Washington, New Hampshire

In addition to national map layers covering percent tree canopy cover in 12-digit HUCs and land cover in stream and lake buffers, EnviroAtlas also features maps for select communities that quantify the amount of pollutants removed, carbon stored, and annual runoff reduced by tree cover.

How can I use this information?

This map layer was derived using the NLCD Percent Tree Canopy dataset. Percent tree canopy records the proportion of land surface that is covered by tree tops within every 30 m² pixel. This approach differs from a land cover classification that classifies an entire 30 m² pixel as forest or other land cover type. Tree canopy cover is a useful measure of intermittent tree distribution in agricultural and urban areas and in riparian buffers.

The map layer, Percent Tree Canopy Cover, may be used to evaluate the potential value of ecosystem services provided by tree canopy. The map indicates which 12-digit HUCs may benefit from additional tree cover. This information can be used to determine areas with a high or low potential for wildlife habitat connectivity, carbon storage, air and water pollutant reduction, and flood storage.

The density and extent of tree cover can be more thoroughly investigated in EnviroAtlas by increasing the transparency of the map over an aerial imagery base map and adding the NLCD 2011 canopy layer data or NHDPlus Flowline data for streams and water bodies. For more information on riparian tree cover, see EnviroAtlas national and community data layers covering stream and lake buffers.

Though this is a national EnviroAtlas map, it may be used with community EnviroAtlas map layers such as removal of pollutants by trees, reduction in annual runoff by tree cover, and tree cover per capita to characterize the extent of tree canopy in the broader landscape surrounding the community.

How were these data created?

These data were generated by using the 2011 National Land Cover Database (NLCD2011) – [Percent Tree Canopy](#) dataset with the [NHDPlusV2](#) Watershed Boundary Dataset (WBD). The mean percent 2011 NLCD tree canopy cover was calculated and recorded for each 12-digit HUC.

What are the limitations of these data?

Though EnviroAtlas uses the best data available, there are limitations associated with the data. The NLCD percent tree canopy data are generated by the United States Forest Service (USFS). The USFS derives canopy cover from both spectral information in Landsat imagery and other available ground and ancillary information and as such there are inherent errors associated with the data. The USFS includes a standard error for every pixel ranging from 0 to 45 percent and a user can investigate these errors by accessing the NLCD data.⁶

The NLCD 2011 percent tree canopy data was not designed to be directly comparable to the NLCD 2001 canopy data. The two data layers were developed using different methods and

sampling densities. Also, the 2001 data defined trees as >5 meters in height, while the 2011 data had no height restrictions, but defined trees by species. Canopy cover is not available for NLCD 2006.

How can I access these data?

EnviroAtlas data can be viewed in the interactive map, accessed through web services, or downloaded. The [NLCD](#) and [NHDPlusV2](#) data are accessible through their respective websites.

Where can I get more information?

A selection of resources related to tree canopy cover is listed below. For additional information on how the data were created, access the [metadata](#) for the data layer from the drop down menu on the interactive map table of contents and click again on metadata at the bottom of the metadata summary page for more details. To ask specific questions about this data layer, please contact the [EnviroAtlas Team](#).

Acknowledgments

EnviroAtlas is a collaborative effort by EPA, its contractors, and project partners. Donald Ebert, EPA, generated the data for this map. Anne Neale, Donald Ebert, and Megan Mehaffey, EPA; David Eskew, EPA Contractor; David J. Nowak, USDA Forest Service; and Sandra Bryce, Woolpert, Inc. contributed to the fact sheet.

Selected Publications

1. Smith, W.H. 1990. [Air pollution and forests](#). Springer-Verlag, New York, New York. 618 p.
2. Nowak, D.J., D.E. Crane, and J.C. Stevens. 2006. [Air pollution removal by urban trees and shrubs in the United States](#). *Urban Forestry and Urban Greening* 4:115–123.
3. Intergovernmental Panel of Climate Change (IPCC). 2014. [Climate Change 2014: Synthesis Report](#). Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
4. Nowak, D.J., and D.E. Crane. 2002. [Carbon storage and sequestration by urban trees in the USA](#). *Environmental Pollution* 116(3):381–389.
5. Nowak, D.J., J. Wang, and T. Endreny. 2007. [Chapter 4: Environmental and economic benefits of preserving forests within urban areas: air and water quality](#). Pages 28–47 in de Brun, C.T.F. (ed.), *The economic benefits of land conservation*. The Trust for Public Land, San Francisco, California.
6. Coulston, J.W., G.G. Moisen, B.T. Wilson, M.V. Finco, W.B. Cohen, and K. Brewer. 2012. [Modeling percent tree canopy cover: A pilot study](#). *Photogrammetric Engineering & Remote Sensing* 78(7):715–727.